



EOM-SF calculation of methylene (CH_2)





EOM-SF:

$$\Psi(M_s=0) = R(M_s=-1)\Psi_0(M_s=1)$$



Ψ_0

Reference



Ψ_i^a

Target states

EOM-SF method yields:


- accurate singlet-triplet gaps;
- accurate geometries and properties;
- describes closed-shell and open-shell low spin states.

Casanova and Krylov, Spin-flip methods in quantum chemistry, PCCP **22** 4326 (2020)



- Import geometry (methylene.xyz) into IQmol
- Use triplet reference (charge=0, multiplicity=3)
- Request "diradical states" of expected symmetry (see frontier orbitals irreps)
- Request S² of CCSD and EOM wavefunctions to check for spin-contamination
- Look at the EOM amplitudes to assign state character.
- **Important energy differences: between the EOM states (not reference-EOM).**

```
$rem
METHOD          = EOM-CCSD
SF_STATES        = [2,0,0,2] ! Request 4 transitions: A1 and B2 irreps
BASIS           = 6-31G
CC_EOM_PROP_TE  = 1      ! Compute S2 of EOM states
CC_REF_PROP_TE  = 1      ! Compute S2 of CCSD reference
GUI             = 2
$end
```



Character table for C_{2v} point group

	E	C ₂ (z)	$\sigma_v(xz)$	$\sigma_v(yz)$	linear, rotations	quadratic
A ₁	1	1	1	1	z	x ² , y ² , z ²
A ₂	1	1	-1	-1	R _z	xy
B ₁	1	-1	1	-1	x, R _y	xz
B ₂	1	-1	-1	1	y, R _x	yz

Product table for C_{2v} point group

	A ₁	A ₂	B ₁	B ₂
A ₁	A ₁	A ₂	B ₁	B ₂
A ₂	A ₂	A ₁	B ₂	B ₁
B ₁	B ₁	B ₂	A ₁	A ₂
B ₂	B ₂	B ₁	A ₂	A ₁

For each EOM-SF state you should get the following sections in the output

1) Excitation energies and orbital description

EOMSF transition 1/A1

Total energy = -38.97364719 a.u. Excitation energy = 0.0086 eV.

R1^2 = 0.9887 R2^2 = 0.0113 Res^2 = 2.06e-06

Conv-d = yes

Amplitude	Transitions between orbitals			
-0.6827	1 (B2) A	->	1 (B2) B	
0.6698	3 (A1) A	->	3 (A1) B	
-0.1569	1 (B2) A	->	2 (B2) B	
-0.1452	2 (A1) A	->	3 (A1) B	
-0.1126	3 (A1) A	->	5 (A1) B	

Summary of significant orbitals:

Number	Type	Irrep	Energy
2	Occ Alpha	2 (A1)	-0.9633
4	Occ Alpha	3 (A1)	-0.4731
5	Occ Alpha	1 (B2)	-0.4116
4	Vir Beta	3 (A1)	0.1363
9	Vir Beta	5 (A1)	0.8458
5	Vir Beta	1 (B2)	0.1735
10	Vir Beta	2 (B2)	0.8981

2) State properties and S²

EOMSF-CCSD transition 1/A1

S² calculation will be performed in double precision

Excited state properties for EOMSF-CCSD transition 1/A1

Dipole moment (a.u.): 0.251378 (X 0.000000, Y 0.000000, Z -0.251378)

R-squared (a.u.): 23.272158 (XX 10.587846, YY 5.830180, ZZ 6.854132)


Gauge origin (a.u.): (0.000000, 0.000000, 0.000000)

Angular momentum (a.u.) against gauge origin:

(X 0.000000i, Y 0.000000i, Z 0.000000i)

Traces of the OPDMs: Tr(AA) 4.000000, Tr(BB) 4.000000

<S²> = 2.000864



Transition irrep.	EOM-SF state	Energy (eV)	S ²	Multiplicity	E _{ex} (eV)
1/A ₁	1/B ₂	0.0086	2.00	Triplet	0.0000
1/B ₂	1/A ₁	1.1777	0.00	Singlet	1.1691
2/A ₁	2/B ₂	2.2171	0.00	Singlet	2.2085
2/B ₂	2/A ₁	4.3607	0.00	Singlet	4.3521